NYSNA Recommendations and Resources for Reopening during COVID-19

Intensive Care Settings: NYSNAs Nursing Education & Practice and Health & Safety



of care on an ICU unit based on clinical needs. Prioritize services that if deferred are most likely to result in patient harm. Establish designated zones where patients can be screened for COVID – 19. Establish designated zones where patients can be screened for COVID – 19. Establish designated zones where patients can be screened for COVID – 19. Establish designated zones where patients can be screened for COVID – 19. Establish designated zones where patients can be screened for COVID – 19. Establish designated zones where patients can be screened for COVID – 19. Establish designated zones where patients can be screened for COVID – 19. Establish designated zones where patients can be screened for COVID – 19. Establish designated zones where patients can be screened for COVID – 19. Establish designated zones where patients can be screened for COVID – 19. Establish designated zones where patients can be screened for COVID – 19. Establish designated zones where patients can be screened for COVID – 19. Establish designated zones where patients can be screened for COVID – 19. Establish designated zones where patients can be screened for COVID – 19. Establish designated zones where patients can be screened for COVID – 19. Establish designated as Non-COVID – vseritized for testing. Maintenance of 6 feet of more distancing between patients. Air handling systems with appropriate directionality, filtration, and exchange rate should be exhausted directly to the outside or through a HEPA filter before recirculation. Establish designated as Non-COVID – vseritized for testing. Maintenance of 6 feet of more distancing between patients. Air handling systems with appropriate directionality, filtration, and exchange rate should be exhausted directly to the outside or through a HEPA filter before recirculation. Establish designated as Non-COVID – vseritized for testing. Maintenance of 6 feet of more distancing between patients. Covilo – Visitor should be wearing N95 or contacts not adequate). Elou and d	Unit	Pre-Screening	Criteria for Testing	Unit Modifications	PPE	Cleaning	Visitor Access	RN to Patient Ratio
For patients undergoing a procedure or operation, a viral test should be performed 24 hours prior to the Por the patient: Surgical masks Departments should start to incorporate elactomeric	Intensive	Evaluate necessity of care on an ICU unit based on clinical needs. Prioritize services that if deferred are most likely to result in patient harm. Establish designated zones where patients can be screened for	PUI's should be prioritized for testing. If patient presents with any of the following they should be tested: Cough Sore throat Chills Repeated shaking with chills Muscle pain Headache Loss of taste or smell Contact with someone known to have COVID-19 Fever of 100 F Shortness of breath For patients undergoing a procedure or operation, a viral test	ICU Units should be designated as Non-COVID –vs-PUI/COVID units. Maintenance of 6 feet of more distancing between patients. Air handling systems with appropriate directionality, filtration, and exchange rate should be installed and maintained. AIRR should have a minimum of 6 air exchanges per hour and air from these rooms should be exhausted directly to the outside or through a HEPA filter before recirculation. All doors should remain closed except when entering or leaving. Create more AIIRs, permanent or temporary. Create negative air pressure for entire ICU areas. Reduce exposure time	All patient care staff should be wearing N95 or better equipment with eye protection (personal glasses or contacts not adequate). Gloves, isolation gown for nurses on COVID units. For Non-COVID units. For Non-COVID units surgical mask, gown, gloves, as per usually and ordinary standards of practice. For the patient: Surgical masks Departments should start to	Hand hygiene by using ABHR with 60- 95% alcohol or washing hands with soap and water for at least 20 seconds. Use EPA approved emerging viral pathogen cleansers. Clean and disinfect all surfaces before and after all treatments. Sanitize all high touch surfaces. Disposable medical supplies should be discarded not	Refuse visitors who have symptoms or test positive on screening. Visitors should be made aware of visitor policy and requirements including wearing surgical masks, handwashing and any limitations on visitation. Limit visiting hours and use of	Patient Ratio RN's should not float between COVID and non-COVID units. Staff should be continuously and routinely screened for symptoms of COVID.



NYSNA Recommendations and Resources for Reopening during COVID-19

Intensive Care Settings: NYSNAs Nursing Education & Practice and Health & Safety



for 14 days prior to the	PAPRs available for certain	the crisis.		
procedure.	tasks.			
	Concurrently:			
	 Conduct air flow inspections to determine what improvements can be made. Begin investigating the use of dedicated downward flow ventilation directly over 			
	each patient area. 3. Air changes should be boosted above 6 per hour, incorporating more outdoor air.			
	4. Conduct air flow inspections to determine problems and arrive at recommendations for moving contaminated air quickly away from patients and staff.			

https://www.cms.gov/files/document/covid-recommendations-reopening-facilities-provide-non-emergent-care.pdfhttps://psnet.ahrq.gov/web-mm/nurse-staffing-ratios-crucible-money-policy-research-and-patient-care nursingeducation.lww.com/blog.entry.html/2016/11/10/the_importance_ofth-GCAE.html

