

NYSNA Recommendations and Resources for Reopening during COVID-19
Perioperative Settings: NYSNAs Nursing Education & Practice and Health & Safety Guidelines



Unit	Pre-Screening	Criteria for Testing	Unit Modifications	PPE	Cleaning	Visitor Access	RN to Patient Ratio
Peri-operative	Consider use of telemedicine for prescreening data collection.	Exposure to someone diagnosed with COVID-19 in the past 14 days; or,	Patients to be kept in isolation room until transferred to the OR.	Because false-negatives may occur with testing, droplet precaution	Surface disinfectants (quaternary ammonium compound with alcohol), ultra violet light, alcohol-based hand rubs on IV pole, double glove, maintain clean and dirty areas, decolonize patients using pre-procedural chlorhexidine wipes, two doses of nasal povidone iodine within one hour of incision and chlorhexidine mouth rinse.	Implement social distancing policy for visitors that meets the then-current local and national recommendations for community isolation practices.	1 (one) Perioperative RN circulator per patient present during entire intraoperative timeline
	<p>CDC recommended test-based strategy</p> <ul style="list-style-type: none"> i. Resolution of fever without the use of fever-reducing medications; and, ii. Improvement in respiratory symptoms; and, iii. Negative results from two SARS-CoV-2 tests \geq 24 hours apart <p>CDC non-test based strategy</p> <ul style="list-style-type: none"> i. At least 72 hours since resolution of fever without the use of fever-reducing medications and improvement in respiratory symptoms; and, ii. At least 7 days since symptoms first appeared. 	Unexplained fever, cough, shortness of breath, chills, muscle pain, headache, sore throat, and/or new loss of taste or smell within the prior two weeks.	<p>For OR procedure, negative pressure isolation room with minimum of 12 air exchanges per hour.</p> <p>Temporary conversion of otherwise positive pressure OR's into negative pressure with independent provision for air conditioning and humidification.</p> <p>Two adjacent OR's with one being utilized for the procedure and the other as anteroom for staff to wear and dispose of all PPE.</p>	<p>(Surgical mask and eye covering) should be used by OR staff for operative cases.</p> <p>Before performing an aerosol-generating procedure, health care providers within the room should wear an N95 mask, eye protection, gloves and a gown.</p>			
<p>Resources:</p> <p>https://www.asahq.org/about-asa/newsroom/news-releases/2020/04/asa-and-apsf-joint-statement-on-perioperative-testing-for-the-covid-19-virus</p> <p>https://www.asahq.org/about-asa/newsroom/news-releases/2020/04/joint-statement-on-elective-surgery-after-covid-19-pandemic</p> <p>https://pubmed.ncbi.nlm.nih.gov/32217947/</p> <p>https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7189905/</p>							

NYSNA Recommendations and Resources for Reopening during COVID-19
Perioperative Settings: NYSNAs Nursing Education & Practice and Health & Safety Guidelines

